

AD-A103 287 VIRGINIA MASON RESEARCH CENTER SEATTLE WASH  
CHEMICAL MESSENGERS OF INFLAMMATION.(U)  
APR 81 J C HOUCK

F/G 6/5

N00014-77-C-0262

NL

UNCLASSIFIED

1-1  
AD-A103 287



END

DATE

FILED

10-81

DTIC

AD A103287

B5  
HC FILE COPY

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

LEVEL 13

## REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS  
BEFORE COMPLETING FORM

1. REPORT NUMBER 000 1AV	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) CHEMICAL MESSENGERS OF INFLAMMATION A054172		5. TYPE OF REPORT & PERIOD COVERED ANNUAL 4/1/80-3/31/81
7. AUTHOR(s) John C. Houck Ph.D., Director, Principal Investigator		8. CONTRACT OR GRANT NUMBER(s) N00014-77-C-0262
9. PERFORMING ORGANIZATION NAME AND ADDRESS Virginia Mason Research Center 1000 Seneca Street, Seattle, WA 98101		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 206-009 1271
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research, Arlington, Virginia 22217		12. REPORT DATE Apr 1981
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 5
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) -		
18. SUPPLEMENTARY NOTES -		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) -		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The purpose of this contract is to support pilot studies into the possibility of utilizing various biological molecules obtained as natural products which can control the immunologic aspects of inflammation (i.e. inhibit lymphocyte function) and thereby possibly prolong the survival of organ grafts after transplantation.		

DTIC  
ELECTE  
AUG 25 1981  
S D A

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-LF-014-660181 8 3894 024  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

13

ANNUAL REPORT

OFFICE OF NAVAL RESEARCH

CONTRACT NO.

N00014-77-C-0262 Mod. #202-134

INVESTIGATOR:

John C. Houck, Ph.D.  
Director  
Virginia Mason Research Center  
1000 Seneca Street  
Seattle, Washington 98101


Date of Project:


April 1, 1980 through March 31, 1981

TITLE:

Chemical Messengers of Inflammation

29 April 1981

  
John C. Houck, Ph.D., Principal Investigator

Accession For	
A&I	<input checked="" type="checkbox"/>
T&B	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Communication	
Ex.	
Distribution/	
Availability Codes	
Avail and/or	
Dist Special	
	

This document has been approved  
for public release and sale; its  
distribution is unlimited.

81 8 24 024

The purpose of this contract is to support pilot studies into the possibility of utilizing various biological molecules obtained as natural products which can control the immunologic aspects of inflammation (i.e. inhibit lymphocyte function) and thereby possibly prolong the survival of organ grafts after transplantation.

This contract is divided, then, into three sub-projects each of which will be summarized succinctly, but separately.

1. Lymphocyte chalone.

For close to two decades it has been postulated that there exist naturally-occurring materials within cells which inhibit specifically the mitotic activity of those cells. These specific endogenous mitotic inhibitors have been given the name "chalone" from a Greek nautical expression meaning "to slack off" the mainsail and slow down. No chalones have, as yet, been unequivocally purified.

We have approached this project and problem by preparing aqueous extracts of fresh calf thymus and isolating the various inhibiting fractions therein of two-way Mixed Lymphocyte Culture produced transformation of lymphocytes in vitro. The assay involves the use of splenic lymphocytes prepared from inbred mice with a histoincompatible loci on their lymphocytes (BALB 6 and C57 BL). These lymphocytes will, in Microwell plates, each transform the other, thus producing large numbers of large vascularized lymphoblasts which can be identified morphologically in tissue culture microscopes (even by biochemists!) and after 72 hours, quantitative determination of this transformation can be made by measuring the six hour incorporation of radioactive thymidine into these cells.

Aqueous extracts of thymus were resolved on ultragel G-200 Sephadex columns into large, middle and small molecular weight components. The large molecular weight material was found to be capable of inhibiting lymphocyte transformation by a cytotoxic process leading to the discovery of Houck's law, namely "Dead cells don't divide". The middle size molecules (molecular weight 40,000 to 60,000 daltons) turned out to apparently inhibit thymidine uptake but did not inhibit lymphoblast transformation morphologically, by its activity on thymidine (i.e. it was a thymidine kinase rendering the label impossible of entrance into the transforming cell) and the smaller molecular weight material contained inhibition of MLC transformation which appeared neither cytotoxic nor due to thymidine kinase activity.

The small molecular weight material, in a range from 10,000 daltons down, was further subdivided on G-25 exclusion column chromatography and 85% of the activity was found at a molecular weight equivalent to bacitracin (i.e. 1400 daltons). Bleomycin and streptomycin sulfate were also standards which indicated the molecular weight of 1400 daltons for the inhibitor. This material was isolated by preparative electrophoresis and identified ultimately as spermine. This polyamine will inhibit lymphocytes at a few micrograms/ml in the presence of fetal calf serum. The calf serum contains an enzyme (polyamine oxidase) which presumably converts the spermine to an aldehyde which, in turn, is capable of inhibiting lymphocyte transformation perhaps five times more effectively on a dose basis than it inhibits the proliferation of fibroblasts in vitro.

A careful perusal of the literature of chalones indicates that a very large number of inhibitory principles could be spermine, either traveling on an anionic polyelectrolyte, like transfer RNA, or on other anionic macromolecules. Spermine has a molecular weight of 200, but because of its very high cationic quality, it binds a sufficiently large amount of water to make a large hole in the solvent which would cause it to be excluded from Sephadex or Biogel with a spuriously high molecular weight of 1400 daltons. Most of the lymphocyte chalone reports in the literature are due largely to spermine. The proof of this is simply to run the two-way MLC studies not in fetal calf serum, but in human serum, which does not contain significant amounts of polyamine oxidase. It takes over 50 times as much spermine to inhibit MLC transformation in human serum than in fetal calf serum.

Careful study using G-25 columns of the material held way back in a small molecular weight area on Ultragel of thymic extracts indicated that another 10% of the inhibitory capacity of this extract was due to cold thymidine nucleotides diluting out the pool size of the thymidine label. The remaining 5% of the inhibitory activity was neither nucleotide nor polyamine and was investigated in the hope that this would prove to be the elusive chalone. The assays of this inhibitory activity were done in human serum.

The final procedure for the isolation of this latter material is as follows: ammonium acetate (50 mmol) extracts of fresh thymus were prepared, and after clarification by centrifugation, precipitated with 60% ethanol. The supernatant from this precipitate was then mixed with large volumes of acetone and a precipitate produced. All of the inhibitor described above was found in this acetate precipitate.

This precipitate was reconstituted in ammonium acetate and subjected to ion-exchange chromatography using either cationic exchanger AG-50 or anionic exchanger AG-1. All of the inhibitory activity passed effectively through the cationic exchanger indicating that the inhibitor was either anionic or neutral in charge and all of it was retained on the anionic inhibitor suggesting that it must, in fact, be an anionic molecule. This material elutes appropriate standardized G-25 column at a molecular weight of 600 daltons.

The inhibitory activity is destroyed by 4N hydrochloric acid hydrolysis (unlike polyamine) and by acetylation indicating a free amino group despite its anionic nature.

This material can be further purified by elution from LH-20 which completes its desalting and produces a peptide fraction which at 20 µg/ml will inhibit by 50% the thymidine uptake of two-way MLC. This fraction, which is still impure, does not inhibit the thymidine uptake of K562 erythroblastoid cells, thus indicating a considerable degree of specificity even at a still impure state.

We are currently investigating the inhibitory capacity of this fraction against B cells (mouse lymphocytes transformed by LPS) and are attempting to further purify it by silicic acid and ultimately HPLC.

## 2. Bacterial Inhibitor

We have cultivated a pure culture of *Pseudomonas* recovered from a patient with a remarkable tolerance for cadaver kidney graft. We have isolated from within this bacterium a protein with a molecular weight of perhaps 80,000 daltons which apparently paralyzes both human and mouse lymphocytes in vitro.

This material is prepared by disrupting the harvested *Pseudomonas aeruginosa* cells by shaking with glass beads at high speed and clarified with streptomycin sulfate to remove the nucleic acids, leaving a clear supernatant. This supernatant is precipitated with 30% alcohol to remove some extraneous components and all of the inhibitory activity is concentrated at 60% precipitate with ethanol. This precipitate is re-dissolved in saline and chromatographed on Ultragel 200, and separated from various proteases which can destroy it. The 80,000 dalton sized portions of the eluate are, in turn, concentrated and subjected to Red Dye Ligand Chromatography. All of the inhibitory activity is eluted at the beginning of the salt gradient from this column and while heterogeneous, still represents highly purified inhibitor fraction.

Preliminary experiments in mice indicate that this material has lost most of its toxicity and, when administered every other day at a dose of 0.1 or 1 mg/mouse, will significantly prolong the survival of C-57 skin transplant on BALB-C mice. Specifically, the normal time required to reject the skin grafts is 10 to 12 days. We can prolong this up to 15 days with the bacterial product at the moment. We are in the process of making more of this material and trying to complete its purification while in more detailed experiments attempting to demonstrate its activity biologically as an immunosuppressive agent.

### 3. Poly-polyamine.

Most recently we have been attempting to develop a series of cross-linked polymers of spermine to study the ability of these increasing-sized polymers to inhibit lymphocyte transformation in vitro. We have prepared a dimer of spermine by cross-linking it with glutaraldehyde and separating the molecule product as a dimer by exclusion column chromatography. Spermine elutes on Sephadex with a molecular weight of apparently 1400 daltons or 7 times its true mass. The dimer, not surprisingly, has an apparent molecular weight of 3500 daltons. Five µg/ml will inhibit lymphocyte transformation in human serum. This is the same activity displayed in fetal calf serum so obviously this dimer does not require or react with polyamine oxidase.

We are going to attempt to develop further the polymerization of polyamine by isolating dimers which will then be further cross-linked with glutaraldehyde. Theoretically, we would hope this might become an increasingly effective molecule for the inhibition of lymphocyte function both in vitro and in vivo.

Preliminary studies have indicated that the polyamine, when dimerized, will, in fact, delay skin graft rejection across histoincompatible mice (as described above) from 12 to 14 days. This is borderline significance, but then the injection of 100 µg of the material per mouse every other day may not be the ideal dose, route or compound (i.e. the tetramer might be better). We hope to explore this during the coming year.

### Summary:

We have been exploring the efficacy of three different types of compounds capable of inhibiting lymphocyte function in vitro in terms of their potential application to control inflammation and particularly, organ graft rejection in vivo. Three papers have been published describing the results of this work to date:

- a) Patt, L.M. and J.C. Houck. The incredible shrinking chalone. Febs. Lett. 120(2): 163-170, Nov. 1980.
- b) Patt, L.M., J.M. Gleisner, D.M. Barrantes and J.C. Houck. Low molecular weight inhibitors of lymphocyte transformation. (In press, Pharmacology).
- c) Gleisner, J.M., L.M. Patt, C.A. Ramthun and J.C. Houck. Pulmonary polyamine permeability factor. Inflammation (In Press).

OFFICE OF NAVAL RESEARCH  
BIOLOGICAL SCIENCES DIVISION  
BIOPHYSICS PROGRAM, CODE 444  
DISTRIBUTION LIST FOR TECHNICAL, ANNUAL AND FINAL REPORTS

# Copies

- (13) Administrator, Defense Documentation Ctr  
Cameron Station  
Alexandria, Virginia 22314
- (6) Director, Naval Research Laboratory  
Attention: Technical Information Div.  
Code 2627  
Washington, D.C. 20375
- (6) Office of Naval Research  
Attention: Code 102IP (ONRL DOC)  
800 N. Quincy St.  
Arlington, Virginia 22217
- (3) Office of Naval REsearch  
Biophysics Program  
Code 444  
Arlington, Virginia 22217
- (1) Commanding Officer  
Naval Medical Res. & Develop. Command  
National Naval Medical Center  
Bethesda, Maryland 20014
- (1) Chief, Bureau of Medicine & Surgery  
Department of the Navy  
Washington, D.C. 20375
- (2) Technical Reference Library  
Naval Medical Research Institute  
National Naval Medical Center  
Bethesda, Maryland 20014
- (1) Office of Naval Research Branch Office  
Building 114, Section D  
666 Summer Street  
Boston, Massachusetts 02210
- (1) Office of Naval Research Branch Office  
536 South Clark Street  
Chicago, Illinois 60605
- (1) Department of the Army  
U.S. Army Science & Technology Ctr. -  
Far East  
APO San Francisco 96328
- (1) Assistant Chief for Technology  
Office of Naval Research, Code 200  
800 North Quincy Street  
Arlington, Virginia 22217

# Copies

- (1) Office of Naval Research Branch Office  
1030 East Green Street  
Pasadena, California 91106
- (1) Commanding Officer  
Naval Medical Research Unit No. 2  
Box 14, APO San Francisco 96263
- (1) Commanding Officer  
Naval Medical Research Unit No. 3  
FPO New York 09527
- (1) Officer in Charge  
Submarine Medical Research Laboratory  
Naval Submarine Base, New London  
Groton, Connecticut 06342
- (1) Scientific Library  
Naval Aerospace Medical Research Institute  
Naval Aerospace Medical Center  
Pensacola, Florida 32512
- (1) Commanding Officer  
Naval Air Development Center  
Attn: Aerospace Medical Res. Dept.  
Warminster, Pennsylvania 18974
- (1) Director  
Naval Biosciences Laboratory  
Building 844  
Naval Supply Center  
Oakland, California 94625
- (1) Commander, Army Research Office  
P.O. Box 12211  
Research Triangle Park  
North Carolina 27709
- (1) Directorate of Life Sciences  
Air Force Office of Scientific Research  
Bolling Air Force Base  
Washington, D.C. 20332
- (1) Commanding General  
Army Medical Research and Develop. Comm  
Forrestal Building, WA D.C. 20314



**DATE**  
**ILME**